

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Immunology and Respiratory Medicine

Journal homepage: <https://www.ijirm.org/>

Original Research Article

Prevalence of hypothyroidism among patients with obstructive sleep apnea: A study in a tertiary care center

M Ravindranath¹, D.S Sowjanya^{1,*}¹Dept, of Pulmonary Medicine, Kamineni Academy of Medical Science & Research Center, Hyderabad, Telangana, India

ARTICLE INFO

Article history:

Received 06-06-2021

Accepted 01-08-2021

Available online 29-09-2021

Keywords:

Obstructive sleep apnea

hypothyroidism

apneahypoapnea index

ABSTRACT

Background: Obstructive sleep apnea is a common disorder wherein there are recurrent episodes of sleep disordered breathing resulting in disrupted sleep and other sequelae. The symptoms of hypothyroidism are very similar to OSA, a number of studies have been performed to analyze their association with conflicting results.

Materials and Methods: A questionnaire based on the Wisconsin Sleep Apnea questionnaire for quality of sleep, sleep pattern, symptoms during sleep, snoring were asked to 118 patients included in the study. The day time sleep patterns were assessed by the Epworth Sleepiness scale. Polysomnography testing, Saturated oxygen levels, sleep efficiency and arousal index were also done for all the patients. Blood was collected for the detection of thyroid hormones.

Results: There were 82 (69.5%) males and 36 (30.5%) females on the present study out of 118 patients. The mean age of the patients under study was 53.91 ± 4.69 years, the weight was 89.48 ± 12.83 kgs and the BMI was 34.86 ± 6.11 . Amongst the patients with OSA and hypothyroid, the apnea hypoapnea index was 66.22 ± 18.31 episodes per hour, and Epworth sleepiness scale was 15.18 ± 5.26 while in patients with OSA and without hypothyroid, it was 24.17 ± 6.29 and 8.25 ± 5.82 respectively.

Conclusions: Our study shows a significant association between OSA and hypothyroidism among patients, thus having a considerable implications in managing the thyroid condition of the patients.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Obstructive sleep apnea is a common disorder wherein there are recurrent episodes of sleep disordered breathing resulting in disrupted sleep and other sequelae.^{1–3} These episodes maybe due to total or partial pharyngeal collapse as well as in temporary obstruction of the airways during sleep thus resulting in hypoxemia and hypocapnia. These symptoms are restored during the arousal of the individual as the pharynx is opened restoring the airflow, but in turn it disturbs the quality of sleep. Many times, this results in excessive sleep during the day, the condition

called as obstructive sleep apnea syndrome (OSAS).¹ Other symptoms associated with the fragmented sleep are apathy, somnolence and lethargy.

OSA is very often accompanied by lowered blood saturation levels. This is due to the pauses in the breath which may last for 20–40 seconds. The patient is normally unaware of these pauses, but it is noticed by the people around him. OSA is characterized by snoring, gasping and choking.^{4–8} Some of the risk factors for OSA are age, obesity, a familial history and the circumference of the neck. OSA is also said to occur more in males in comparison to the female population. In case of the females, menopause is said to be the main cause.^{9–11}

* Corresponding author.

E-mail address: sowjanya@yahoo.in (D. S. Sowjanya).

Since the symptoms of hypothyroidism are very similar to OSA, a number of studies have been performed to analyze their association with conflicting results. The mechanisms associated for OSA have been proposed to be an increased mucopolysaccharide and protein deposition in the upper respiratory system with an alteration in the regulatory control of the dilator muscles of the pharynx due to neuropathy and a depression of the respiratory centers.^{12,13}

According to National Health and Nutrition Examination Survey (NHANES) done in the USA, the prevalence of hypothyroidism was estimated to be around 3.7%, while sleep apnea was around 3%. Using different criteria, the prevalence of hypothyroidism among the patients with OSA has been reported to be around 1-10%.^{14,15}

Since OSA is a risk factor for many diseases such as cardiovascular diseases, depression, obesity, diabetes, hypertension, the diagnosis of this condition is very important for an effective treatment.^{16,17} It has also been reported that in patients with hypothyroidism, the frequency of OSA is increased.¹⁸ This study was therefore conducted to study the association between OSA and hypothyroidism.

2. Materials and Methods

The present study was conducted by the Department of Pulmonary Medicine at Kamineni Academy of Medical Sciences and Research Centre during the period of two years between December 2019 to March 2021 on 118 patients who were diagnosed with having obstructive sleep apnea. This study was conducted after obtaining clearance for the Institutional Ethical Committee and informed consent from the concerned patients or their family members. In case of minors, the consent was obtained from the parents or the legal guardians. All the patients were subjected to a thorough clinical examinations by the clinician during the initial screening procedure. A questionnaire based on the Wisconsin Sleep Apnea questionnaire was prepared and questions regarding the quality of sleep, sleep pattern, symptoms during sleep, snoring were asked to the patients. The day time sleep patterns were assessed by the Epworth Sleepiness scale.

Polysomnography testing was done for all the patients. For electronic data, the scoring system was done manually as per the established criteria. The sleep apnea in patient was diagnosed as per the definition of International Classification of Sleep Disorders and classification was as per the American Academy of Sleep Medicine (AASM) criteria (Table 1).

Table 1: Classification of OSA as per American Academy of Sleep Medicine (AASM)

Apnea – Hypopnea Index	Criteria of OSA
1-14/h	Mild
15-29 /h	Moderate
≥ 30/h	Severe

Saturated oxygen levels, sleep efficiency and arousal index were also done for all the patients and noted.

Blood was collected from the medial cubital vein within 2 weeks of the sleep apnea test for serum thyroxine (T4) and thyroid stimulating hormone (TSH). The hypothyroidism was divided as following¹⁹ (Table 2)

Table 2: Classification of hypothyroidism

TSH and T4 level	Classification
TSH >5.0 μ IU/mL & T4 <10.3 pmol/L	Clinical hypothyroidism
TSH > 5.0 μ IU/mL & T4 normal range	Subclinical hypothyroidism

3. Results

There were 82 (69.5%) males and 36 (30.5%) females on the present study out of 118 patients (Figure 1).




Fig. 1: Gender classification of patients

The mean age of the patients under study was 53.91 ± 4.69 years, the weight was 89.48 ± 12.83 kgs and the BMI was 34.86 ± 6.11 . Most of the patients in the study were either overweight or obese (Table 3).

Table 3: Demographic detail of the patients

Details	Mean \pm SD
Age (in years)	53.91 ± 4.69
Weight (in Kg)	89.48 ± 12.83
BMI (in kg/m^2)	34.86 ± 6.11

103 (87.3%) of the patients had a history of snoring. Most of the times, this was noticed by the patients attenders. Apnea was observed in 101 (85.6%) of the patients, 78 (66.1%) had somnolence during the day and out of the 118 patients, 22 (18.6%) had hypothyroid (Table 4).

Amongst the patients with OSA and hypothyroid, the apnea hypoapnea index was 66.22 ± 18.31 episodes per

Table 4: Clinical Features of patients

Details	Number (%)
History of snoring	103 (87.3%)
Apnea	101 (85.6%)
Somnolence during day	78 (66.1%)
Hypothyroidism	22 (18.6%)

hour, while in patients with OSA and without hypothyroid, it was 24.17 ± 6.29 . The Epworth Sleepiness scale was 15.18 ± 5.26 in OSA patients with Hypothyroid and 8.25 ± 5.82 in patients without hypothyroid. The sleep efficiency was 68.49 ± 16.39 and 76.19 ± 13.03 % in OSA patients with hypothyroid and without hypothyroid respectively. Arousal index in the patients was 67.96 ± 21.84 in OSA patients with hypothyroid and 57.66 ± 18.73 in patients without hypothyroid (Table 5).

Table 5: Comparison of clinical symptoms in OSA patients with and without hypothyroid

Features	OSA with hypothyroid	OSA without Hypothyroid
Apnea Hypoapnea Index	66.22 ± 18.31	24.17 ± 6.29
Epworth Sleepiness scale	15.18 ± 5.26	8.25 ± 5.82
Sleep efficiency	68.49 ± 16.39	76.19 ± 13.03
Desaturation Index	44.71 ± 11.56	30.15 ± 7.21
Time of SaO ₂ < 90% (min)	34.88 ± 16.44	26.33 ± 13.81
Arousal Index	67.96 ± 21.84	57.66 ± 18.73

The most common risk factors among the patients was hypertension as seen in 71(60.2%) of the patients and diabetes was seen in 49(41.5%). 55(46.6%) of the patients had a history of smoking, some with more than 5 packs a day, while depression was observed in 37(31.4%) of the patients. Few patients (31 (26.3%)) had asthma and 24(20.3%) had a history of coronary heart disease (Table 6).

Table 6: Risk factors

Risk factors	Number	Percentage
Diabetes mellitus	49	41.5%
Hypertension	71	60.2%
Depression	37	31.4%
Smoking	55	46.6%
Asthma	31	26.3%
Coronary arterial Disease	24	20.3%

4. Discussion

Thyroid disorders and obstructive sleep apnea are two disorders commonly found in adult population. OSA is said

to be more associated with obesity among men in whom the disorder is noted to be twice more common than in women¹. Moreover, OSA is also associated with metabolic disorders such as obesity, acromegaly apart from hypothyroidism. It is also suggested that OSA is one of the risk factors for cardiovascular disease.

In the present study, 69.5% of the patient with OSA were males and 30.5% were females. In another study by Young et al, 9% of women and 24% men showed sleep disordered breathing, with day time somnolence.¹

The mean age was 53.91 ± 4.69 years with a mean BMI of $34.86 \pm 6.11 \text{ kg/m}^2$. In a study by BaHannan et al., the mean age of the patients in their study was 48.7 ± 14.1 years²⁰ with a BMI of $37.7 \pm 9.6 \text{ kg/m}^2$ corroborating our study. The prevalence of apnea in obese patients is probably due to the deposits of fat in the pharynx lateral wall reducing the caliber of UAs.²¹ Obesity was one of the main factors for sleep apnea. In a study by Kapur et al., Women aged below 50 years were found to be more prone to OSA.²²

83.7% of the patients in the present study had a history of snoring, while apnea was observed in 85.6% of the patients. 66.1% of the patients showed somnolence during the day. Hypothyroidism was seen in 18.6% of the patients. In a study by BaHammam et al., the prevalence of hypothyroidism was 11.5%²⁰ and was more prevalent in the female group. In several studies, an increase in the development of OSA is observed in patients with hypothyroidism.^{23,24} 11% of hypothyroidism was diagnosed in OSA patients in another study by Popovivi et al.,²⁵

Amongst the patients with OSA and hypothyroid, the apnea hypoapnea index was 66.22 ± 18.31 episodes per hour, while in patients with OSA and without hypothyroid, it was 24.17 ± 6.29 . In a similar study by Rajgopal et al., a range of 17 -176 episodes per hour were seen among the patients.²⁶ They also observed obese patients to have considerably more episodes of apnea compared to the non-obese patients.²¹ In a similar study by BaHammam et al., the AHI was 55.2 ± 37 /hr in OSA group of patients.⁹

The ESS scale was 15.18 ± 5.26 in patients with hypothyroid and 8.25 ± 5.82 in without hypothyroid. There was not much difference between the two groups in the arousal index. The time of SaO₂ < 90% /min. Similar were the cases in studies by BaHammam et al., and Miller et al.,^{9,27}

In studies by Skjodt et al, the patients who were treated with thyroxine were found to have improved response to sleep apnea.²⁸ Similar results were observed in other studies by Millman et al and Orr et al while Skatrud et al., and Grunstein et al., found no difference in sleep apnea after treatment with thyroxine.^{13,29-31}

5. Conclusions

Our study shows a significant association between OSA and hypothyroidism among patients, thus having a considerable implications in managing the thyroid condition of the patients. Thus, we recommend routine thyroid estimation in patients with OSA, so that if detected, it can be treated at the earliest.

6. Acknowledgments

None.

7. Conflict of interests

The authors declare that there are no conflicts of interest in this paper

8. Source of Funding

None

References

- Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S, et al. The occurrence of sleep-disordered breathing among middle aged adults. *N Engl J Med.* 1993;328(17):1230–5.
- Lawati NMA, Patel SR, Ayas NT. Epidemiology, risk factors and consequences of obstructive sleep apnea and short sleep duration. *Prog Cardiovasc Dis.* 2009;51(4):285–93. doi:10.1016/j.pcad.2008.08.001.
- Remmers JE, Degroot WJ, Sauerland EK, Anch AM. Pathogenesis of upper airway occlusion during sleep. *J Appl Physiol.* 1978;44(6):931–8. doi:10.1152/jappl.1978.44.6.931.
- Obstructive Sleep Apnea Syndrome (780.53-0). The international classification of sleep disorders. Westchester, Illinois: American Academy of Sleep Medicine; 2001. p. 52–60.
- Bresnitz EA, Goldberg R, Kosinski RM. Epidemiology of obstructive sleep apnea. *Epidemiol Rev.* 1994;16(2):210–27. doi:10.1093/oxfordjournals.epirev.a036151.
- Millman RP, Fogel BS, Mcnamara ME, Carlisle CC. Depression as a manifestation of obstructive sleep apnea: reversal with nasal continuous positive airway pressure. *J Clin Psychiatry.* 1989;50(9):348–51.
- Findley LJ, Levinson MP, Bonnier RJ. Driving performance and automobile accidents in patients with sleep apnea. *Clin Chest Med.* 1992;13(3):427–35.
- Madani M, Madani F. Epidemiology, pathophysiology, and clinical features of obstructive sleep apnea. *Oral Maxillofac Surg Clin North Am.* 2009;21(4):369–75. doi:10.1016/j.coms.2009.09.003.
- Bahammam AS, Alrajeh MS, Al-Jahdali HH, Binsaeed AA. Prevalence of symptoms and risk of sleep apnea in middle-aged Saudi males in primary care. *Saudi Med J.* 2008;29(3):423–9.
- Tsara V, Amfilochiou A, Papagrigorakis JM, Georgopoulos D, Liolios E, Kadiths A, et al. Guidelines for diagnosing and treating sleep related breathing disorders in adults and children (Part 3: obstructive sleep apnea in children, diagnosis and treatment). *Hippokratia.* 2010;14(1):57–62.
- Ancoli-Israel S, Klauber MR, Kripke DF, Parker L, Cobarrubias M. Sleep apnea in female patients in a nursing home. *Chest.* 1989;96(5):1054–8. doi:10.1378/chest.96.5.1054.
- Kapur VK, Koepsell TD, Demaine J, Hert R, Sandblom RE, Psaty BM, et al. Association of hypothyroid and obstructive sleep apnea. *Am J Respir Crit Care Med.* 1998;158(5 Pt 1):1379–83. doi:10.1164/ajrccm.158.5.9712069.
- Gruntsein RR, Sullivan CE. Sleep apnea and hypothyroidism: mechanism and management. *Am J Med.* 1988;85(6):775–9.
- Winkelman JW, Goldman H, Piscatelli N, Lukas SE, Dorsey CMS. Cunningham Are thyroid function tests necessary in patients with suspected sleep apnea? *Sleep.* 1996;19(10):790–3.
- Grunstein R. Obstructive sleep apnea syndrome and hypothyroidism. *Chest.* 1994;105(4):1296–7.
- Pinto JA, Ribeiro DK, Cavallini AF, Duarte C, Freitas GS. Comorbidities associated with obstructive sleep apnea: a retrospective study. *Int Arch Otorhinolaryngo.* 2016;20(2):145–50.
- Sweed RA, Hassan, Elwahad HA, Aref SR, Mahmoud MI. Comorbidities associated with obstructive sleep apnea: a retrospective Egyptian study on 244 patients. *Sleep Breath.* 2019;23(4):1079–85. doi:10.1007/s11325-019-01783-w.
- Mete T, Yalcin Y, Berker D. Relationship between obstructive sleep apnea syndrome and thyroid diseases. *Endocrine.* 2013;44(3):723–8. doi:10.1007/s12020-013-9927-9.
- Surks MI, Ortiz E, Daniels GH, Sawin CT, Col NF, Cobin RH, et al. Subclinical thyroid disease: scientific review and guidelines for diagnosis and management. *JAMA.* 2004;291(2):228–38.
- Bahammam A, Sharif M, Jammah A, Bahammam S. Prevalence of Thyroid disease in patients with Obstructive Sleep Apnea. *Respir Med.* 2011;105(11):1755–60. doi:10.1016/j.rmed.2011.07.007.
- Shelton KE, Woodson H, Gays S, Suratt PM. Pharyngeal fat in obstructive sleep apnea. *Am Rev Respir Dis.* 1993;148(2):462–6. doi:10.1164/ajrccm/148.2.462.
- Vishesh KK, Thomas DK, Demaine J, Hert R, Robert ES, Bruce MP, et al. Association of Hypothyroidism and Obstructive Sleep Apnea. *Am J Respir Crit Care Med.* 1998;158(5):1379–83. doi:10.1164/ajrccm.158.5.9712069.
- Hira HS, Sibal L. Sleep apnea syndrome among patients with hypothyroidism. *J Assoc Physicians India.* 1999;47(6):15–8.
- Jha A, Sharma SK, Tendon N, Lakshmy R, Kadhiraavan T, Handa KK, et al. Thyroxine replacement therapy reverses sleep disordered breathing in patients with [primary hypothyroidism. *Sleep.* 2006;7(1):55–61. doi:10.1016/j.sleep.2005.05.003.
- Popovici I, Khawaja I. Efficacy of thyroid function tests in patients suspected of having obstructive sleep apnea. *Chest.* 1997;112:149.
- Rajagopal KR, Abbrecht PH, Derderian SS, Pickett C, Hofeldt F, Tellis CJ, et al. Obstructive sleep apnea in hypothyroidism. *Ann Intern Med.* 1984;101(4):491–4.
- Miller CM, Husain AM. Should women with obstructive sleep apnea syndrome be screened for hypothyroidism? *Sleep Breath.* 2003;7(4):185–8.
- Skjodt NM, Atkar R, Easton PA. Screening for hypothyroidism in sleep apnea. *Am J Respir Crit Care Med.* 1999;160(2):732–5.
- Millman RP, Bevilacqua J, Peterson DD, Pack AI. Central sleep apnea in hypothyroidism. *Am Rev Respir Dis.* 1983;127(4):504–7. doi:10.1164/arrd.1983.127.4.504.
- Orr WC, Males JL, Imes NK. Myxedema and obstructive sleep apnea. *Am J Med.* 1981;70(5):1061–6. doi:10.1016/0002-9343(81)90867-6.
- Skatrud J, Iber C, Ewart R, Thomas G, Rasmussen H, Schultze B, et al. Disordered breathing during sleep in hypothyroidism. *Am Rev Respir Dis.* 1981;124(3):325–9. doi:10.1164/arrd.1981.124.3.325.

Author biography

M Ravindranath, Associate Professor

D.S Sowjanya, Assistant Professor

Cite this article: Ravindranath M, Sowjanya DS. Prevalence of hypothyroidism among patients with obstructive sleep apnea: A study in a tertiary care center. *IP Indian J Immunol Respir Med* 2021;6(3):161-164.