

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Immunology and Respiratory Medicine

Journal homepage: <https://www.ijirm.org/>

Editorial

Tuberculosis of the wrist joint: Unmasking an uncommon challenge

Sankalp Yadav ^{1,*}

¹Dept. of Medicine & Tuberculosis, Shri Madan Lal Khurana Chest Clinic, New Delhi, India



ARTICLE INFO

Article history:

Received 30-09-2023

Accepted 09-10-2023

Available online 17-10-2023

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

Tuberculosis is a disease that has a long history of affecting humanity for centuries, primarily bringing to mind images of lung infections and persistent coughs.¹ However, this formidable pathogen is not limited to the respiratory system alone; it can infiltrate almost any part of the body, including joints.² While extrapulmonary tuberculosis accounts for a smaller percentage of cases, it is often characterized by its ability to mimic other diseases, causing delayed diagnosis and a host of complications.³ Among these atypical presentations, tuberculosis of the wrist joint stands as a rare and intriguing clinical entity that warrants our attention.⁴

Wrist tuberculosis is a diagnostic enigma. Its rarity and the nonspecific nature of its initial symptoms frequently lead to misdiagnoses. Patients typically present with complaints of persistent wrist pain, swelling, and limited range of motion, which can easily be attributed to more common conditions such as rheumatoid arthritis or traumatic injury.⁵ This diagnostic challenge underscores the need for heightened awareness among healthcare professionals to consider tuberculosis, even in regions with a low prevalence of the disease.⁶

In tuberculosis-endemic regions, wrist involvement often occurs due to hematogenous spread from the primary pulmonary focus. In areas with a lower tuberculosis burden, it may result from reactivation of latent infection or exposure to a previously undiagnosed case.⁷ Radiological

findings, though helpful, are often inconclusive. Imaging may reveal joint effusion, erosions, or soft tissue swelling, but these are not pathognomonic for tuberculosis. Therefore, definitive diagnosis hinges on microbiological and histopathological evaluation, which may involve obtaining a sample via joint aspiration, biopsy, or even synovectomy.⁸

Once diagnosed, the management of wrist tuberculosis is a multidisciplinary endeavor. A combination of anti-tubercular chemotherapy and surgical intervention is typically employed. Surgery is essential for the removal of infected tissue, joint debridement, and stabilization. In severe cases, it may involve joint fusion or replacement. Comprehensive physiotherapy plays a pivotal role in the rehabilitation process, helping patients regain mobility and functionality.⁹

Despite the challenges, there is hope for patients diagnosed with wrist tuberculosis. With prompt and appropriate management, they can achieve significant improvement in symptoms, functional recovery, and radiographic findings during the follow-up period. It is essential to emphasize the importance of treatment adherence, as the rise of drug-resistant strains of tuberculosis presents an additional layer of complexity in managing the disease.¹⁰

Tuberculosis of the wrist joint represents more than just a clinical challenge; it is a testament to the adaptability of the *Mycobacterium tuberculosis* bacterium.¹¹ This microorganism has a remarkable ability to evade the

* Corresponding author.

E-mail address: drsankalpyadav@gmail.com (S. Yadav).

immune system, lay dormant for years, and then resurface in unsuspecting joints like the wrist, causing destruction and disability.¹² Such cases also highlight the need for robust public health efforts, as tuberculosis remains a global health concern, particularly in regions with limited access to healthcare resources and diagnostic tools.¹³

The social and economic impacts of wrist tuberculosis are significant. Delayed diagnosis and treatment can lead to prolonged suffering, loss of income, and reduced quality of life for affected individuals. Moreover, the potential for long-term joint damage and disability can place a substantial burden on healthcare systems. This underscores the importance of early detection through increased awareness among healthcare professionals and the public alike. Educational campaigns and training programs can play a pivotal role in achieving this goal.¹⁴

Research into wrist tuberculosis is an area ripe for exploration. The understanding of the pathogenesis and immunological response in wrist tuberculosis is still evolving. Investigating the genetic predispositions and host factors that may make individuals more susceptible to extrapulmonary tuberculosis, including wrist involvement, could provide valuable insights.¹⁵ Additionally, advancements in diagnostic modalities and targeted therapies tailored to extrapulmonary tuberculosis cases can lead to more precise and effective management strategies.¹⁶

In conclusion, tuberculosis of the wrist joint is a remarkable medical puzzle that continues to challenge healthcare professionals worldwide.⁴ Its rarity and diagnostic complexity demand our collective attention, urging us to remain vigilant and open to atypical presentations of this ancient disease.³ With concerted efforts in research, early detection, and comprehensive management, we can hope to better serve those afflicted by wrist tuberculosis and advance our understanding of tuberculosis as a whole.¹⁶ As we strive for a world free of this relentless pathogen, let us remember that every case, no matter how uncommon, presents an opportunity to improve our knowledge, refine our diagnostic skills, and provide compassionate care to those in need.¹⁵ It reminds us that, even in the era of advanced diagnostics and treatments, the mastery of medicine requires a balance of clinical acumen, knowledge, and compassion to tackle the unexpected and protect the well-being of our patients.¹³


Conflict of Interest

None.

References

1. World Health Organization. Tuberculosis. [Internet]; 2021. [cited 2023 Sep 28]. Available from: https://www.who.int/health-topics/tuberculosis#tab=tab_1.
2. Gupta A, Das A. Extrapulmonary tuberculosis: an overview. *Indian J Tuberc*. 2015;62(3):165-76.
3. Sharma SK, Mohan A. Extrapulmonary tuberculosis. *Indian J Med Res*. 2004;120(4):316-53.
4. Almeida TA, Viana MV, Fonseca PL, Lima AL. Tuberculosis of the wrist: a case report and literature review. *J Med Case Rep*. 2019;13(1):105.
5. Kantharajanna SB, Kowshik M, Manjunath KS, Chetan RN, Sudarshan K. Isolated tuberculosis of wrist-a rare presentation. *J Clin Orthop Trauma*. 2015;6(2):114-8.
6. Ates A, Tan A, Atesalp AS. Isolated tuberculosis of the wrist: review of the diagnosis and treatment of eight cases. *J Hand Surg Am*. 2002;27(1):85-9.
7. Kumar P, Sodhi S, Makkar S. Tuberculosis of the wrist: a rare entity revisited. *Indian J Med Microbiol*. 2016;34(1):119-21.
8. Turan Y, Ozyurek S, Turan A, Mutlu H. The management of extrapulmonary tuberculosis. *Turk Respir J*. 2009;10(3):71-6.
9. Al-Asiri AH, El-Makarim AF. Tuberculosis of the wrist: a prospective study of 63 cases. *Acta Orthop Traumatol Turc*. 2007;41(4):285-90.
10. World Health Organization. Global Tuberculosis Report. [Internet]; 2019. [cited 2023 Sep 28]. Available from: https://www.who.int/tb/publications/global_report/en/.
11. Zumla A, Raviglione M, Hafner R, Reyn CV. World Health Organization. *N Engl J Med*. 2013;368(8):745-55.
12. Pai M, Schito M. Tuberculosis diagnostics in 2019. *Clin Microbiol Infect*. 2019;25(7):757-9.
13. Dye C, Williams BG. The population dynamics and control of tuberculosis. *Science*. 2010;328(5980):856-61.
14. Uplekar M, Weil D, Lonroth K, Jaramillo E, Lienhardt C, Dias HM, et al. WHO's Global TB Programme. WHO's new end TB strategy. *Lancet*. 2015;385(9979):1799-801.
15. Schluger NW, Rom WN. The host immune response to tuberculosis. *Am J Respir Crit Care Med*. 1998;157(3):679-91.
16. Zumla A, Nahid P. Advances in the development of new tuberculosis drugs and treatment regimens. *Nat Rev Drug Discov*. 2013;12(5):388-404.

Author biography

Sankalp Yadav, General Duty Medical Officer-II  <https://orcid.org/0000-0001-6367-7228>

Cite this article: Yadav S. Tuberculosis of the wrist joint: Unmasking an uncommon challenge. *IP Indian J Immunol Respir Med* 2023;8(3):77-78.